MCT PLASMA

Breaking PRP limits

BEAUTYFORM MEDICAL



In the last decade,
improving PRP
solely on
This limit has now
MCT Plasma moves one
dramatically increasing
and enhancing

performance depended
platelet concentration.
been overcome.
step beyond PRP,
platelet performance
its results.

MCT PLASMA
Much more than PRP

MCT is a medical system that improves autologous products such as PRP or stem cells concentrates. It delivers all the power of regenerative medicine in a very easy and clever way. By managing electromagnetic and thermal energy, **MCT system transforms traditional PRP into MCT Plasma.**

MCT Plasma is a new generation autologous product that stands out for its much higher growth factor concentration, remarkable ATP production and spectacular skin rejuvenation, tissue regeneration and aesthetic results.

MCT PLASMA

STIMULATES CELLS AND PRP PERFORMANCE

ATP

X2 ATP PRODUCTION

x3

INCREASES GROWTH FACTORS

MCT System

MCT System is the result of 3 years of R&D aiming to create a device capable of bringing the benefits of Photo-Thermal activation to medical consultations. Its results are impressive and its potential breathtaking.

- Enhanced Regenerative power
- Stunning platelet performance
- More comfortable injection
- Extraordinary clinical and aesthetic results

MCT Unit®

MCT Unit® enables existing regenerative medicine treatments to be boosted, simply and quickly. Through the application of electromagnetic and thermal energy, it improves the general characteristics of autologous products, such as growth factor content, ATP production, and cell or platelet performance. It handles a wide range of temperatures and multiple wavelengths which are set up in comfortable one-touch protocols and customized programs.

MCT Kit®

MCT Kit® is a patented device with a specific shape and chemical composition, exclusively developed for photothermal conditioning. It is made of a medical grade polymer that guarantees optimal scattering, transmitance, and other optical properties, ensuring that the emitted light will reach the target effectively. Its unique geometry grants an excellent surface/volume energy exposure ratio. MCT Kit® can allocate 10 mL of any autologous preparate.



3
easy steps to get MCT Plasma

Step 01

Procure the autologous material

Step 02

Pour it into the MCT Kit®

Step 03

Set up the MCT Unit®

MCT Plasma
Breaking PRP limits

Much more than PRP...

Increment of Growth Factors

Photo-Thermal boosted platelets release a higher amount of growth factors in a controlled and sustained way for a more physiological effect.

Improved Platelet Performance

Photo-Thermal stimulated plasma contains less proinflammatory cytokines and has a more physiological pH. It also improves platelet activation profile, reducing VASP and increasing p-38 phosphorilation. Cold primes platelets and facilitates its reactivity once injected.

Reduction of Pain

Cold primes alpha granules release kinetics after activation, not before, and induces a delay in clot formation. A less dense plasma and the sensation the cold liquid produces result in a much more comfortable procedure.

ATP Synthesis Stimulation

Photo-Thermal boosting stimulates cellular metabolism and enhances ATP synthesis.

PRP PHOTO BOOSTER

Light provides a non-invasive, non-toxic way of stimulating cellular metabolism. Ultimately, the energy absorbed at the mitochondria enhances cellular ATP production. Photo-Thermal stimulation of PRP generates MCT Plasma, a state-of-the-art regenerative preparate. This vanguard autologous plasma achieves superb clinical results due to a more prolonged and sustained growth factors release and a reduced pro-inflammatory cytokine content. Its remarkable results can be seen not only in the rejuvenation of face, neck and hands, but also in the awesome hair growth witnessed in patients with androgenic alopecia.



Fvasconcellos 22:35, 9 September 2007 (UTC),
Public domain, via Wikimedia Commons

Total control of the energy



Biphasic dose response in LLLT. Three-dimensional plot illustrating effects of varying irradiation time equivalent to fluence or irradiance on the biological response resulting in stimulation or inhibition.

If the light applied is not of sufficient irradiance or the irradiation time is too short, then there will be no response. If the irradiance is too high or irradiation time is too long, then the response may be inhibited. Somewhere in between is the optimal combination of irradiance and time for stimulation.

Text and figure extracted from Chung H, Dai T, Sharma SK, Huang YY, Carroll JD, Hamblin MR. The nuts and bolts of low-level laser (light) therapy. Ann Biomed Eng. 2012 Feb;40(2):516-33. doi: 10.1007/s10439-011-0454-7. Epub 2011 Nov 2. PMID: 22045511; PMCID: PMC3288797.

PRP THERMAL BOOSTER

Temperature has a huge body of evidence that backs up its effectiveness and safe utilization. Thermal stimulation prepares platelets to work more and better, increases their reactiveness and shows-up a myriad of activation markers. Platelet primed functionality also determines a huge increment in the release of growth factors. Since thermal boosting delays the clot onset and primes platelet aggregation only after activation, MCT Plasma results in a less dense regenerative product, implying not only the achievement of spectacular clinical results but also a much less painful injection experience.



Extracted from Kobsar A, Koehnlechner K, Klingler P, Niklaus M, Zeller-Hahn J, Koessler A, Weber K, Boeck M, Koessler J. The effect of short-term refrigeration on platelet responsiveness. Sci Rep. 2022 Oct 7;12(1):16910. doi: 10.1038/s41598-022-21124-4. PMID: 36207457; PMCID: PMC9546855. Adapted from figure 8, created with BioRender.com Mechanism of enhanced platelet reactivity by refrigeration. Attenuated inhibitory signaling facilitates ADP- and collagen-induced aggregation and platelet adhesión or aggregate formation.

Energy control at every level

GROWTH FACTORS



Comparison of mean concentrations of 3 growth factors measured in activated PRP before (blue) and after (orange) thermal conditioning at 4°C for 10 minutes. Concentrations before (pre-TC) and after (post-TC) thermal conditioning are described as means and standard deviations. EGF: pre-TC 21.98 (3.41), post-TC 43.69 (4.05). p-value<0.001. Basic FGF: pre-TC 9.188 (1.94), post-TC 16.981 (2.57). p-value<0.001. VEGF: pre-TC 4.88 (0.89) and post-TC 6.99 (0.79). p-value<0.001.

Extracted from Melamed G, et al. Thermal Conditioning: im provingPRP growth factro content. J Prime 2020 june;2:2-5”

MCT PLASMA PRIMES PLATELET PERFORMANCE FOR AWESOME CLINICAL AND AESTHETIC RESULTS



Youthful Hands





Ageless Décolleté

M·C·T

10 minutes of PRP photo-conditioning increases 2-3-folded growth factors and ATP



- Light enables sustained growth factor release from PRP.
- Significant ATP increment ($p<0.001$) in all groups after photostimulation.
- Photoactivated platelets have significantly higher ($p<0.001$) percentage of P-selectin expression as compared resting platelets.

Cumulative release (pg/uL) of growth fators in PRP: A – PDGF, B – TGF-ß, C – bFGF. Comparison of ATP secretion release from unstimulated and photostimulated platelets (D). Data are expressed as mean value of triplicates and error bars as the standard deviation. Photo Stimulation 1 min (PL-PAC1), Photo Stimulation 5 min (PL-PAC5), Photo Stimulation 10 min (PL-PAC10).
Extracted from G. Irmak, T.T. Demirtaş, M. Gümü şderelioğlu, Sustained Release of Growth Factors from Photoactivated Platelet Rich Plasma (PRP), European Journal of Pharmaceutics and Biophar- maceutics (2019), doi: https://doi.org/10.1016/j.ejpb.2019.11.011.

4°C Thermal stimulated PRP shows more physiologic charateristics



t-PRP demonstrated natural fiber scaffolding, which trapped more platelet and GFs, and exhibited a slow release and degradation rate of GFs.

t-PRP is a novel and convenient method for the preparation and activation of PRP without any additives. Compared to contemporary PRP, t-PRP reflects more physiologic characteristics.

ELISA protein quantification of GF of t-PRP and c-PRP groups. Total accumulation of GFs over 0.5–96 h for VEGF and PDGF-AB.
Extracted from Du L, Miao Y, Li X, Shi P, Hu Z. A Novel and Convenient Method for the Preparation and Activation of PRP without Any Additives: Temperature Controlled PRP. Biomed Res Int. 2018 May 13;2018:1761865. doi:10.1155/2018/ 1761865. PMID: 29862255; PMCID: PMC5971324.

Release of VEGF, EGF and bFGF is significantly increased in 4°C thermo-conditioned PRP



PRP is usually processed at 37°C or room temperature (RT 20–25°C); however, it is traditionally known that lower temperatures prime platelet activation and trigger the release of alpha granules. The release of VEGF, EGF, bFGF was significantly increased when PRP was incubated at 4°C before inducing coagulation.

Cold preconditioning promotes the release of growth factors and cytokines from platelets. (A) PRP was incubated at 37 °C, 23 °C, or 4 °C for 30 min, and the levels of VEGF, EGF, bFGF and PDGF in releasates of PRP before clotting (CaCl2−) or after clotting (CaCl2+) were determined by ELISA. Total intra-platelet levels were measured in platelet lysates.
Extracted from fig. 2, Etulain J, Mena HA, Meiss RP, Frechtel G, Gutt S, Negrotto S, Schattner M. An optimised protocol for platelet-rich plasma preparation to improve its angiogenic and regenerative properties. Sci Rep. 2018 Jan 24;8(1):1513. doi: 10.1038/s41598-018-19419- 6. PMID: 29367608; PMCID: PMC5784112.

Efficacy and safety of photothermal-bioactivated platelet-rich plasma for facial rejuvenation



Compared to standard PRP, Photothermal-bioactivated PRP showed excellent results for facial rejuvenation.

Extracted from figure 1, Beltrán B, Sánchez MAR, Melamed G, Pinto H. Efficacy and safety of photothermal-bioactivated platelet-rich plasma for facial rejuvenation. J Cosmet Dermatol. 2022 Jul 20. doi: 10.1111/jocd.15250. Epub ahead of print. PMID: 35855670.

Courtesy: Dr. Manuel Rubio, 2022. CIME Clinic, Barcelona.

Beltrán B et al, 2022. Efficacy and safety of photothermal-bioactivated platelet-rich plasma for facial rejuvenation. J Cosmet Dermatol. 2022 Jul 20. doi: 10.1111/jocd.15250. Epub ahead of print. PMID: 35855670.

Hernández C, et al. Efficacy of photothermoactivated PRP for skin biostimulation in patients not eligible for other medical-aesthetic treatments. Publication 2023.

Tejero P, et al. Impact of photothermoactivated PRP on skin rejuvenation of the hands. Publication 2023.

R&D

Efficacy and Safety of Photo-thermal-bioactivated PRP for Facial Rejuvenation.

Beltrán B, Sánchez MAR, Melamed G, Pinto H. J Cosmet Dermatol. 2022 Jul 20. doi:10.1111/jocd.15250.

Effect of Photobiomodulation on PRP: Review Series on New Tools in Regenerative Medicine.

Pinto H, Sánchez-Vizcaíno Mengual E. J Aest Med. 2021;7(4):31-7

The Effect of Photobiomodulation on Human Mesenchymal Cells: A Literature Review.

Pinto H, Goñi Oliver P, Sánchez-Vizcaíno Mengual E. Aest Plast Surg. 2021Aug;45(4):1826-42

Thermal Conditioning of PRP: improving growth factor content.

Pinto H, Melamed G. J Prime. 2020june;2:2-5

Photoactivation of Autologous Materials with a New Reliable, Safe and Effective Set-Up.

Pinto H. J Aest Med. 2020; 6(1):11-5.

Efficacy of photothermoactivated PRP for skin biostimulation in patients not eligible for other medical-aesthetic treatments.

Hernández C, et al. Publication 2023.

Impact of photothermoactivated PRP on skin rejuvenation of the hands.

Tejero P, et al. Publication 2023.

Photo-Thermal stimulated PRP (MCT PLASMA) provides excellent results for hair restoration. A 150-patient experience.

Mercuri S et al, 2023. San Rafaele Hospital, Milan.

Inspiring technology.
Outstanding results.

MCT
PRP
BAPAMs
CUSTOM

Leading edge technology
Evidence-based
Less painful
Safe
Fast
User friendly

Bibliografía

1. G. Irmak, T.T. Demirtaş, M. Gümü şderelioğlu, Sustained Release of Growth Factors from Photoactivated Platelet Rich Plasma (PRP), European Journal of Pharmaceutics and Biopharmaceutics (2019), doi: https://doi.org/10.1016/j.ejpb.2019.11.011

2. Etulain J, Mena HA, Meiss RP, Frechtel G, Gutt S, Negrotto S, Schattner M. An optimised protocol for platelet-rich plasma preparation to improve its angiogenic and regenerative properties. Sci Rep. 2018 Jan 24;8(1):1513. doi: 10.1038/s41598-018-19419-6. PMID: 29367608; PMCID: PMC5784112.

3. Du L, Miao Y, Li X, Shi P, Hu Z. A Novel and Convenient Method for the Preparation and Activation of PRP without Any Additives: Temperature Controlled PRP. Biomed Res Int. 2018 May 13;2018:1761865. doi: 10.1155/2018/1761865. PMID: 29862255; PMCID: PMC5971324.

4. Pinto H, Goñi Oliver P, Sánchez-Vizcaíno Mengual E. The Effect of Photobiomodulation on Human Mesenchymal Cells: A Literature Review. Aesthetic Plast Surg. 2021 Aug;45(4):1826-1842. doi: 10.1007/s00266-021-02173-y. Epub 2021 Feb 22. PMID: 33616715.

5. Beltrán B, Sánchez MAR, Melamed G, Pinto H. Efficacy and safety of photothermal-bioactivated platelet-rich plasma for facial rejuvenation. J Cosmet Dermatol. 2022 Jul 20. doi: 10.1111/jocd.15250. Epub ahead of print. PMID: 35855670.

6. Chung H, Dai T, Sharma SK, Huang YY, Carroll JD, Hamblin MR. The nuts and bolts of low-level laser (light) therapy. Ann Biomed Eng. 2012 Feb;40(2):516-33. doi: 10.1007/s10439-011-0454-7. Epub 2011 Nov 2. PMID: 22045511; PMCID: PMC3288797.

7. Huang YY, Chen AC, Carroll JD, Hamblin MR. Biphasic dose response in low level light therapy. Dose Response. 2009 Sep 1;7(4):358-83. doi: 10.2203/dose-response.09-027.Hamblin. PMID: 20011653; PMCID: PMC2790317.

8. Kim WS, Calderhead RG. Is light-emitting diode phototherapy (LED-LLLT) really effective? Laser Ther. 2011;20(3):205-15. doi: 10.5978/islsm.20.205. PMID: 24155530; PMCID: PMC3799034.

9. Pinto H. Photoactivation of Autologous Materials with a New, Reliable, Safe and Effective Set-Up. J Aest Med. 2020; 6(1):11-5.

10. Kobsar A, Koehnlechner K, Klingler P, Niklaus M, Zeller-Hahn J, Koessler A, Weber K, Boeck M, Koessler J. The effect of short-term refrigeration on platelet responsiveness. Sci Rep. 2022 Oct 7;12(1):16910. doi: 10.1038/s41598-022-21124-4. PMID: 36207457; PMCID: PMC9546855.

11. Melamed G, Pinto H. Thermal Conditioning: improving growth factor content. Prime-Journal. 2020june;2:2-5.

12. Karu T. Primary and secondary mechanisms of action of visible to near-IR radiation on cells. J Photochem Photobiol B. 1999 Mar;49(1):1-17. doi: 10.1016/S1011-1344(98)00219-X. PMID: 10365442.

13. Karu TI, Kolyakov SF. Exact action spectra for cellular responses relevant to phototherapy. Photomed Laser Surg. 2005 Aug;23(4):355-61. doi: 10.1089/pho.2005.23.355. PMID: 16144476.

14. Pinto H, Sánchez-Vizcaíno Mengual E. Effect of Photobiomodulation on PRP: Review Series on New Tools in Regenerative Medicine. J Aest Med. 2021;7(4):31-7.

15. Ferraresi C, Kaippert B, Avci P, Huang YY, de Sousa MV, Bagnato VS, Parizotto NA, Hamblin MR. Low-level laser (light) therapy increases mitochondrial membrane potential and ATP synthesis in C2C12 myotubes with a peak response at 3-6 h. Photochem Photobiol. 2015 Mar-Apr;91(2):411-6. doi: 10.1111/php.12397. Epub 2014 Dec 30. PMID: 25443662; PMCID: PMC4355185.

16. Tafur J, Mills PJ. Low-intensity light therapy: exploring the role of redox mechanisms. Photomed Laser Surg. 2008 Aug;26(4):323-8. doi: 10.1089/pho.2007.2184. PMID: 18665762; PMCID: PMC2996814.

17. Jeyaraman M, Muthu S, Jeyaraman N, Gupta A. Photoactivated platelet-rich plasma: is it the future of platelet-rich plasma? Regen Med. 2022 Sep;17(9):607-609. doi: 10.2217/rme-2022-0063. Epub 2022 Jun 15. PMID: 35703033.

18. Flores Luna GL, de Andrade ALM, Brassolatti P, Bossini PS, Anibal FF, Parizotto NA, Leal ÂMO. Biphasic Dose/Response of Photobiomodulation Therapy on Culture of Human Fibroblasts. Photobiomodul Photomed Laser Surg. 2020 Jul;38(7):413-418. doi: 10.1089/photob.2019.4729. Epub 2020 Mar 24. PMID: 32208063.

19. Akiyama M, Takami H, Yoshida Y. The mechanism of cold-induced platelet aggregation in the presence of heparin. Tohoku J Exp Med. 1995 Dec;177(4):365-74. doi: 10.1620/tjem.177.365. PMID: 8928196.

20. Hoffmeister KM, Felbinger TW, Falet H, Denis CV, Bergmeier W, Mayadas TN, von Andrian UH, Wagner DD, Stossel TP, Hartwig JH. The clearance mechanism of chilled blood platelets. Cell. 2003 Jan 10;112(1):87-97. doi: 10.1016/s0092-8674(02)01253-9. PMID: 12526796.

21. Maurer-Spurej E, Pfeiler G, Maurer N, Lindner H, Glatter O, Devine DV. Room temperature activates human blood platelets. Lab Invest. 2001 Apr;81(4):581-92. doi: 10.1038/labinvest.3780267. PMID: 11304578.

22. Kim JI, Bae HC, Park HJ, Lee MC, Han HS. Effect of Storage Conditions and Activation on Growth Factor Concentration in Platelet-Rich Plasma. J Orthop Res. 2020 Apr;38(4):777-784. doi: 10.1002/jor.24520. Epub 2019 Nov 19. PMID: 31709604.

23. Egidi MG, D'Alessandro A, Mandarello G, Zolla L. Troubleshooting in platelet storage temperature and new perspectives through proteomics. Blood Transfus. 2010 Jun;8 Suppl 3(Suppl 3):s73-81. doi: 10.2450/2010.012S. PMID: 20606754; PMCID: PMC2897204.

24. Etulain J, Lapponi MJ, Patrucchi SJ, Romaniuk MA, Benzadón R, Klement GL, Negrotto S, Schattner M. Hyperthermia inhibits platelet hemostatic functions and selectively regulates the release of alpha-granule proteins. J Thromb Haemost. 2011 Aug;9(8):1562-71. doi: 10.1111/j.1538-7836.2011.04394.x. PMID: 21649851; PMCID: PMC3155010.

25. ZUCKER MB, BORRELLI J. Reversible alterations in platelet morphology produced by anticoagulants and by cold. Blood. 1954 Jun;9(6):602-8. PMID: 13160109.

26. White JG, Krivit W. An ultrastructural basis for the shape changes induced in platelets by chilling. Blood. 1967 Nov;30(5):625-35. PMID: 6073858.

27. Kattlove HE, Alexander B. The effect of cold on platelets. I. Cold-induced platelet aggregation. Blood. 1971 Jul;38(1):39-48. PMID: 4254206.

M·C·T
Meta Cell Technology

BEAUTYFORM MEDICAL

Exclusive Provider of META CELL TECHNOLOGY, S.L.

Beautyform Medical Ltd.
121 Gloucester Place,
Marylebone, W1U 6JY
London

info@beautyform.co.uk
0330 1220879

CE 0051

ISO 13485:2016

EU-MDR

WIPO WORLD INTELLECTUAL PROPERTY ORGANIZATION

www.beautyform.co.uk